

On the absence of fifth-order contributions to the nucleon mass in heavy-baryon chiral perturbation theory

Judith A. McGovern and Michael C. Birse

*Theoretical Physics Group, Department of Physics and Astronomy
University of Manchester, Manchester, M13 9PL, U.K.*

We have calculated the contribution of order M_π^5 in the chiral expansion of the nucleon mass in two-flavour heavy-baryon chiral perturbation theory. Only one irreducible two-loop integral enters, and this vanishes. All other corrections in the heavy-baryon limit can be absorbed in the physical pion-nucleon coupling constant which enters in the M_π^3 term, and so there are no contributions at M_π^5 . Including finite nucleon mass corrections, the only contribution agrees with the expansion of the relativistic one-loop graph in powers of M_π/m_N , and is only 0.3% of the M_π^3 term. This is an encouraging result for the convergence of two-flavour heavy-baryon chiral perturbation theory.

The fundamental degrees of freedom of the strong interaction are quarks and gluons, but in spite of the many successes of QCD in describing high energy phenomenology, a full description of the particles that constitute ordinary matter still eludes us. There is however increasing interest in the interactions of such particles at low energies as new, precise, data on nucleon properties and interactions becomes available. For low enough energy it has long been known that the interaction of pions is governed only by the symmetries of QCD, in particular $SU(2) \times SU(2)$ chiral symmetry, and a successful systematic effective field theory of pions, Chiral Perturbation Theory [1], has been developed. In this theory the pionic lagrangian is expanded as a power series, with each subsequent term having two more derivatives or powers of the pion mass than the previous one, and with the lowest order being two: $\mathcal{L}_{\pi\pi} = \mathcal{L}_{\pi\pi}^{(2)} + \mathcal{L}_{\pi\pi}^{(4)} + \dots$. The expansion is in powers of q/Λ , where q is a momentum or pion mass, and Λ is the scale of the physics which has not been included explicitly, for instance the ρ meson. Because chiral symmetry requires the pion-pion scattering length to vanish in the chiral limit as the pion momentum goes to zero, loop diagrams with vertices from $\mathcal{L}_{\pi\pi}^{(2)}$ only contribute at fourth order; divergences are cancelled by counterterms in $\mathcal{L}_{\pi\pi}^{(4)}$, and so on. At fourth order eight *a priori* undetermined low energy constants (LEC's) enter, which have to be fit to data. The current state of the art is calculation to sixth order, involving two-loop integrals [2].

Attempts to expand the success of the pionic theory by including nucleons initially ran into difficulties; since the nucleon mass is not small, it spoils the expansion of physical amplitudes in powers of small quantities (masses and momenta), and the theory is not systematic [3]. However if the nucleon mass is taken to be infinite, it effectively decouples, and the power counting is restored. The Lagrangian expansion includes both odd and even numbers of derivatives, $\mathcal{L}_{\pi N} = \mathcal{L}_{\pi N}^{(1)} + \mathcal{L}_{\pi N}^{(2)} + \dots$. Only even powers of the pion mass enter, since the underlying param-

eter is the quark mass; $M_\pi^2 \sim M_q$. Furthermore corrections for a finite mass can be included systematically, with a simultaneous expansion in powers of $1/m_N$; since $m_N \sim \Lambda \sim M_\rho$ these terms can naturally be fitted into the expansion above; thus $\mathcal{L}_{\pi N}^{(2)}$ contains terms of order $1/m_N$, and $\mathcal{L}_{\pi N}^{(3)}$, of order $1/m_N^2$, etc.

The heavy-baryon expansion starts by splitting the nucleon momentum into two parts, $p_\mu = mv_\mu + l_\mu$, with $v^2 = 1$ and $v \cdot l \ll m$. The projection operators $P_v^\pm = (1 \pm \not{v})/2$ are used to split the nucleon spinor into two parts, $\Psi = e^{-imv \cdot x}(H + h)$, and the “small-component” degrees of freedom h are integrated out. For details, the reader should consult, for instance, Ref. [4].

Heavy-baryon chiral perturbation theory (HBCPT) has been applied to a number of problems involving the electromagnetic properties of nucleons and the interaction of nucleons and pions. The number of terms to a given order however is much larger than in the pionic theory; there are for instance seven LEC's in $\mathcal{L}_{\pi N}^{(2)}$ and twenty three in $\mathcal{L}_{\pi N}^{(3)}$. So far the full Lagrangian has only been worked out to third order, though individual terms have been considered at higher order. By no means all of the LEC's to third order have been determined from experiment. Although many of the calculations which have been done show the method to be promising, the convergence of the expansion of amplitudes in powers of the physical pion mass is hard to judge so far. In particular, almost all calculations so far have been to one-loop order only*.

One of the quantities which, through its relative simplicity, lends itself to such an analysis, is the nucleon mass shift due to the finite quark (pion) mass, or equiv-

*Bernard *et al.* [5] consider the contribution of two-loop diagrams to the imaginary part of the nucleon isoscalar electromagnetic formfactor, but this does not require the evaluation of two-loop integrals.



Fig. 1: The one-loop diagram of $\Sigma^{(3)}$. Solid lines are nucleons and dashed lines, pions.

alently the pion-nucleon sigma commutator, $\sigma_{\pi N} = M_\pi^2 \partial m_N / \partial M_\pi^2$, estimated from scattering data to be 45 ± 8 MeV [6]. To order q^3 there are two contributions, with one LEC which has now been estimated independently from pion-nucleon scattering data [8,7]. To order q^4 more LEC's will enter, for which there we as yet have no experimental handle. Borasoy and Meißner have attempted to estimate fourth order LEC's in three flavour HBCPT through the principle of resonance saturation [9]. There are however good reasons to distrust three flavour calculations, at least without an explicit decuplet, since nucleonic excitations which are being kept (such as ΣK) are much higher in energy than others which have been integrated out (the delta and higher resonances) [10]. (Convergence of the three-flavour result also appears to be poor [11].) However the mass shift is one quantity where it is possible to look at the order q^5 term without knowing the q^4 piece, and this is what we have done in this paper.

In order to calculate most quantities to order q^5 , the expansion of the nucleonic Lagrangian up to $\mathcal{L}_{\pi N}^{(5)}$ would be required. The only contribution to $\Sigma(0)$ from the fifth order term would be a simple counterterm of order M_π^5 . However the Lagrangian is analytic in the quark masses, that is in M_π^2 , so such a term cannot exist. (For the same reason any irreducible two-loop diagrams must give finite contributions to $\Sigma(0)$, since there can be no counterterm to cancel divergences.) Similarly, in the absence of mass insertions of order M_π , $\mathcal{L}_{\pi N}^{(4)}$ cannot contribute at this order. All the relevant Feynman amplitudes for these calculations can be found in the work of Meißner *et al* [4,7] or, using an alternative form of $\mathcal{L}_{\pi N}^{(3)}$, in that of Mojžiš [8] (who also gives the relevant amplitudes from $\mathcal{L}_{\pi N}^{(4)}$).

The heavy-baryon propagator is given by

$$S^{-1} = \omega - \Sigma(\omega, \mathbf{k}), \quad (1)$$

where the nucleon momentum is written as $p = mv + k$, m is the bare mass, and $\omega = v \cdot k$. The mass shift $\delta m = m_N - m$ is the value of ω for which the propagator has a pole at zero three-momentum:

$$\delta m - \Sigma(\delta m, 0) = 0. \quad (2)$$

(Of course the mass shift could be found from the pole of the propagator for any three-momentum, but as HBCPT is constructed to respect Lorentz invariance, the result will not change.)

In order to solve Eq. 2 to a given order in M_π , both δm and $\Sigma(\omega)$ must be expanded in powers of M_π . To order M_π^3 , $\Sigma(\delta m) = \Sigma(0)$, so [4]

$$\delta m^{(2)} + \delta m^{(3)} = -4c_1 M_\pi^2 - \frac{3g_A^2 M_\pi^3}{32\pi F_\pi^2}, \quad (3)$$

where the second term comes from the diagram in Fig. 1. Writing as $\Sigma^{(n)}$ the expression for the $O(q^n)$ part of Σ , which will have an expansion $\Sigma^{(n)}(\omega) = a_1 M_\pi^n + a_2 M_\pi^{(n-1)} \omega + \dots$, we obtain

$$\delta m^{(5)} = \Sigma^{(5)}(0) + \delta m^{(2)} \Sigma^{(4)'}(0) + \delta m^{(3)} \Sigma^{(3)'}(0) + \frac{1}{2} (\delta m^{(2)})^2 \Sigma^{(3)''}(0) \quad (4)$$

where primes indicate derivatives with respect to ω . Any calculation in HBCPT yields an answer in terms of the bare Lagrangian parameters, M, g, F and m to lowest order, which are the first terms in an expansion in powers of M of M_π, g_A, F_π and m_N . It is customary to replace the bare parameters by the physical ones so that the lowest order predictions do not change as higher orders are added. This however gives an extra contribution to the higher order calculations. In this case, therefore, to the calculation of $\Sigma^{(5)}(0)$ from the diagrams of Fig. 2 must be added a piece from $\Sigma^{(3)}(0)$. Here the relevant parameters are M_π and the pseudo-vector πN coupling $f_{\pi NN} \equiv g_{\pi NN}/m_N = g/F(1 + O(M_\pi^2))$.

The diagrams which contribute to $\Sigma^{(5)}(0)$ are shown in Fig. 2. They consist of two-loop diagrams, and one-loop diagrams with either one insertion from $\mathcal{L}_{\pi N}^{(3)}$ or $\mathcal{L}_{\pi\pi}^{(4)}$, or two insertions from $\mathcal{L}_{\pi N}^{(2)}$. In all cases only zero external momentum is required. The two-loop diagrams shown in Fig. 2f-i all vanish trivially. The diagram in Fig. 2a can after some algebra be written as $(3M^2/4(2d-3)F^4)I$, where

$$I = \int \frac{d^d l d^d k}{(2\pi)^{2d}} \frac{1}{v \cdot k (M^2 - l^2)(M^2 - (k-l)^2)}. \quad (5)$$

This integral can be done by using Feynman parameters, first combining the mesonic propagators in the standard way and integrating over l , and then using the identity

$$\frac{1}{P^p Q^q} = \frac{2^q \Gamma(p+q)}{\Gamma(p)\Gamma(q)} \int_0^\infty \frac{y^{q-1} dy}{(P+2yQ)^{p+q}}$$

to include the heavy-baryon propagator. Integration over y and k yields

$$\begin{aligned} I &= -\frac{M^{2d-5} \pi^{\frac{1}{2}} \Gamma(\frac{5}{2}-d)}{(4\pi)^d} \int_0^1 (x-x^2)^{(1-d)/2} dx \\ &= -\frac{M^{2d-5} 2^{d-2} \pi \Gamma(\frac{5}{2}-d) \Gamma(\frac{3-d}{2})}{(4\pi)^d \Gamma(2-\frac{d}{2})} \end{aligned} \quad (6)$$

which tends to zero as $d \rightarrow 4$.

The same integral also appears in the evaluation of Fig. 2b and 2d, and it is the only non-separable two-loop




Fig. 2: Contributions to $\Sigma^{(5)}$. Solid dots represent insertions from $\mathcal{L}_{\pi N}^{(3)}$ and $\mathcal{L}_{\pi\pi}^{(4)}$, and crosses from $\mathcal{L}_{\pi N}^{(2)}$ (both include fixed terms from the expansion in $1/m_N$).




Fig. 3: Contributions to $\Sigma^{(4)}$.

integral which does. Since it vanishes, all the two-loop diagrams 2a-e are proportional to $\Delta_\pi J_0(0)$, in the notation of [4]. The integral $J_0(0) = -M/8\pi$ is the one which enters the one-loop diagram, Fig. 1, and $\Delta_\pi = 2M^2 L(M)$ is just the integral of the meson propagator and diverges as $1/(d-4)$. These divergences are cancelled by the graphs of Fig. 2j-l with counterterm insertions from $\mathcal{L}_{\pi N}^{(3)}$ and $\mathcal{L}_{\pi\pi}^{(4)}$, which however bring in the low energy constants $2\bar{d}_{16} - \bar{d}_{18}$ from the baryonic, and $\bar{l}_3$ and $\bar{l}_4$ from the mesonic Lagrangians. (The notation is that of [7], with LEC's defined to absorb the usual factors of $\log(M/\mu)$; in [8] $\bar{d}_n \rightarrow \bar{d}_{n+1}$ and $\bar{l}_n \rightarrow \bar{l}_n/16\pi^2$.) The graphs with two insertions from $\mathcal{L}_{\pi N}^{(2)}$, Fig. 2o-q, are all finite. (Counterterm graphs 2m-o all give vanishing contributions.) The final contribution of Fig. 2, with the conventions of [7], is

$$\Sigma_{2\text{-loop+CT}}^{(5)}(0) = \frac{3g^2 M^5}{32\pi F^2} \left(\frac{2\bar{l}_4 - 3\bar{l}_3}{F^2} - \frac{4(2\bar{d}_{16} - \bar{d}_{18})}{g} + \frac{g^2}{32\pi^2 F^2} + \frac{1}{8m^2} + \frac{6c_1}{m} + 24c_1^2 \right). \quad (7)$$

The next contribution to $\delta m^{(5)}$ is from the last three terms of Eq. 4; the relevant graphs are shown in Fig. 3 and Fig. 1, and the integrals are finite. The result is

$$\begin{aligned} & \delta m^{(3)} \Sigma^{(3)'}(0) + \delta m^{(2)} \Sigma^{(4)'}(0) + \frac{1}{2} (\delta m^{(2)})^2 \Sigma^{(3)''}(0) \\ &= \frac{3g^2 M^5}{32\pi F^2} \left(\frac{3g^2}{32\pi^2 F^2} - 4c_1 \left(\frac{3}{2m} + 12c_1 \right) + 24c_1^2 \right). \quad (8) \end{aligned}$$

It may be seen that all terms involving the LEC c_1 vanish from the sum of Eqs (7,8).

Finally we need the contribution to $\Sigma^{(5)}(0)$ from replacing the bare constants in $\Sigma^{(3)}(0)$ with their physical values. (The expression for $\Sigma^{(3)}(0)$ may be found from that for $\delta m^{(3)}$ in Eq. 3 by reinstating the bare coupling constants.) The diagrams which contribute to the renormalised πN coupling $f_{\pi NN} (= g_{\pi NN}/m_N)$ are given in Fig. 4.

The last diagram of Fig. 4 indicates the contribution from the expansion of the pion and nucleon wavefunction renormalisation to order M_π^2 . Some comment is required about Z_N , which is defined as the residue of the nucleon propagator at the pole. This has recently been the subject of a paper by Ecker and Mojžiš [12], who point out a correction which is necessary in order to reproduce the results of a relativistic calculation in HBCPT. It arises because a nucleon can also be created by the eliminated

“small” component of the relativistic nucleon field, and so the normalisations of the relativistic and heavy baryons do not match. The same correction was included through the spinor normalisation by Fearing *et al.* [13]. The net effect of including the small-component sources, to order q^3 , is to give

$$Z_N = (1 + \mathbf{k}^2/4m_N^2)Z_N^{\text{HB}}, \quad (9)$$

where Z_N^{HB} is calculated purely from the HBCPT Lagrangian. (Whereas in the relativistic theory Z_N is a constant, in HBCPT it may depend on the on-shell three-momentum $\mathbf{k}$.) In the framework of Ref. [7], which we have been using here,

$$Z_N^{\text{HB}} = 1 - \mathbf{k}^2/4m_N^2 + \dots, \quad (10)$$

where other terms of order q^2 have been suppressed. Thus in this framework, the dependence on $\mathbf{k}$ cancels to this order.[†] With this small-component-source correction, we find that there are no $O(1/m^2)$ corrections to $f_{\pi NN}$. (In [12] such terms are also shown to be absent from g_A , so the usual expression for the Goldberger-Treiman discrepancy [4,7], proportional only to $\bar{d}_{16}$, holds.) Thus we obtain for the physical pion mass and πN coupling constant,

$$M_\pi^2 = M^2(1 + 2\bar{l}_3 M^2/F^2) \quad (11)$$

$$f_{\pi NN} = \frac{g}{F} \left[1 - M^2 \left(\frac{g^2}{16\pi^2 F^2} + \frac{\bar{l}_4}{F^2} - \frac{4\bar{d}_{16} - 2\bar{d}_{18}}{g} \right) \right]$$

and substituting in $\sigma^{(3)}(0)$ to obtain the final contribution to $\delta m^{(5)}$ gives

$$\Sigma_{1\text{-loop}}^{(5)}(0) = \frac{3g^2 M^5}{32\pi F^2} \left(-\frac{g^2}{8\pi^2 F^2} - \frac{2\bar{l}_4 - 3\bar{l}_3}{F^2} + \frac{4(2\bar{d}_{16} - \bar{d}_{18})}{g} \right). \quad (12)$$

Collecting all contributions, Eq. (7,8,12), we obtain our final result,

$$\delta m^{(3)} + \delta m^{(5)} = -\frac{3g_{\pi NN}^2 M_\pi^3}{32\pi m_N^2} \left(1 - \frac{M_\pi^2}{8m_N^2} \right). \quad (13)$$

Thus in the heavy-baryon limit the order M_π^5 contribution vanishes, with all corrections being absorbed in the physical pion mass and pion-nucleon coupling constant in the M_π^3 contribution. For finite nucleon mass, the correction is just that obtained if the relativistic one-loop contribution is expanded in powers of M_π/m_N

[†]In [12], Z_N in this framework is given wrongly, since the $O(1/m^2)$ kinetic energy insertion, Eq. C1 of [7], has been missed. We have repeated our calculations in Ecker and Mojžiš's framework, and obtained the same final result.




Fig. 4: Contributions to the pion-nucleon vertex of order M_π^2 .

[3]. (Since the $1/m_N$ terms in the HBCPT Lagrangian are constructed to respect Lorentz invariance, this agreement is reassuring but certainly not surprising.)

There remains the question of why there is no other fifth order correction, apart from the renormalisation of the coupling constants of the lowest-order theory and $1/m_N$ corrections. In fact the only thing which could give such corrections would be irreducible two-loop integrals from the diagrams of Fig. 2a, 2b and 2d. As detailed above, only one such integral enters, and in four dimensions it vanishes. This vanishing seems to be essentially accidental. It does not occur in odd dimensions, nor in all probability for unequal meson masses. (We have been able to calculate it for one massless and one massive meson in four dimensions, with a finite but non-zero result.)

Interestingly, it has been conjectured that the main contributions at order q^5 would come from insertions of vertices from $\mathcal{L}_{N\pi}^{2,3}$ in one-loop diagrams and not from genuine two-loop diagrams [9]. Since the two-loop integral vanishes, this is trivially satisfied; however the extent to which other terms are absorbed in the physical coupling constants has not hitherto been realised.

As set out in Eq. 13, the third and fifth order pieces in the expansion of the nucleon mass in powers of M_π are now known. For a full picture of the convergence, we need to know the second and fourth order pieces. The latter have now been calculated, [14] and fall into two categories; pieces involving new LEC's from the fourth order Lagrangian, and fixed pieces. The latter are very small, but the LEC's in the former are not known. As well as entering the nucleon mass, these affect the pion-nucleon sigma term and πN scattering. If c_1 were well determined from scattering, the analysis of which has currently been carried out only to order q^3 , then comparison with the sigma term would tell us whether or not there was still a significant discrepancy to be made up from fourth or higher order contributions. Unfortu-

nately the analyses are not sensitive to c_1 , though the preferred value would leave a substantial M_π^4 contribution. None-the-less it would appear at least that the odd and even power series are converging separately.

-
- [1] J. Gasser and H. Leutwyler, Ann. Phys. (N.Y.) **158** (1984) 142; Nucl. Phys. **B250** (1985) 465.
 - [2] For a review see J. Bijnens, [hep-ph/9710341](#).
 - [3] J. Gasser, M. E. Sainio and A. Švarc, Nucl. Phys. **B307** (1988) 779.
 - [4] V. Bernard, N. Kaiser and U-G. Meißner, Int. J. Mod. Phys. E **4** (1995) 193.
 - [5] V. Bernard, N. Kaiser and U-G. Meißner, Nucl. Phys. **A611** (1996) 429.
 - [6] J. Gasser, H. Leutwyler and M. E. Sainio, Phys. Lett. B **253** (1991) 252, 260.
 - [7] N. Fettes, U-G. Meißner and S. Steininger, Nucl. Phys. **A640** (1998) 199.
 - [8] M. Mojžiš, Eur. Phys. J. C **2** (1998) 181.
 - [9] B. Borasoy and U-G. Meißner, Ann. Phys. (N.Y.) **254** (1997) 19.
 - [10] M. K. Banerjee and J. Milana Phys. Rev. **D52** (1995) 6451.
 - [11] J. F. Donoghue and B. R. Holstein [hep-ph/9803312](#).
 - [12] G. Ecker and M. Mojžiš, Phys. Lett. B **410** (1997) 266.
 - [13] H. W. Fearing, R. Lewis, N. Mobed and S. Scherer, Phys. Rev. **D56** (1997) 1783.
 - [14] S. Steininger, U-G. Meißner and N. Fettes, J. High Energy Phys. **9809** (1998) 8